



# Sero-surveillance of *Mycobacterium avium* subspecies *paratuberculosis* Infection in Domestic Livestock in North India Using Indigenous Absorbed ELISA Test

A.V. Singh, S.V. Singh\*, P.K. Singh, J.S. Sohal and K. Mahour

\*Microbiology Laboratory, Animal Health Division, CIRG, Makhdoom, PO Farah, Mathura- (UP), India.

**Abstract:** A total of 829 serum samples belonging to domestic livestock (cattle, buffaloes, goats and sheep) and driven from different parts of North India between 2005 to 2008, were screened to estimate the seroprevalence of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) infection using 'indigenous absorbed ELISA kit'. The seroprevalence of MAP in domestic livestock was 23.1%. Prevalence was higher in large ruminants (24.1%) as compared to small ruminants (22.5%). The highest seropositivity was in cattle (26.9%), followed by goats (23.9%), buffaloes (20.2%), and sheep (19.0%). In cattle region-wise, 25.8, 29.1 and 30.7% of animals were positive from Mathura (UP), Rohtak (Haryana), and Bareilly (UP) regions, respectively. In buffaloes, the highest prevalence was found at Bareilly (26.6%) followed by Rohtak (20.0%) and Baghpat (18.4%) regions. In goats, 19.6, 37.5, 40.0 and 21.9% of animals were positive from Mathura (farm herd), Etawah, Agra and Ajmer (farmers herd) regions, respectively. In sheep, the prevalence of MAP was 25.5 and 16.3% in Mathura and Mannavanur regions, respectively. In sheep, prevalence was higher in the Northern region as compared to the Southern region of the country. The present study showed that the prevalence of MAP in domestic livestock was moderately higher; therefore there is an urgent need to control the disease at the National level to improve per-animal productivity in the country.

**Keywords:** Johne's disease, Paratuberculosis, *Mycobacterium avium* subspecies *paratuberculosis*, Seroprevalence, ELISA.

## 1. Introduction

*Mycobacterium avium* subsp. *paratuberculosis* (MAP), the cause of Johne's disease (JD) or Paratuberculosis in animals is incurable, chronic wasting and debilitating disease characterized by weight loss and profuse diarrhea (Whitlock and Buergelt, 1996). Though JD is principally a disease of ruminants, MAP can also infect non-ruminant animal species (pigs, dogs, horses, cats, etc.) including free-ranging animals (Blue bull, deer, rabbits) and primates (Chiodini *et al.*, 1984). JD adversely affects animal productivity leading to huge economic losses and has been recognized as one of the most costly infectious diseases of dairy cattle in the world (Hasonova and Pavlik, 2006). The morbidity rates are very high and distributed over a period of time, thus losses in production go unnoticed and have never been estimated in India despite low per-animal productivity (Barbaruah and Joshep, 2008). A countrywide survey of MAP infection has not been undertaken so far, though JD is

endemic in domestic livestock (Singh *et al.*, 2008a; Sharma *et al.*, 2008; Kumar *et al.*, 2007). The country at present lacks any commercial indigenous, cost-effective and sensitive diagnostic kits. Culture through the 'Gold standard test' for JD (Rideout *et al.*, 2003), due to high cost, low sensitivity and the long incubation period, has limited use. PCR cannot distinguish between dead pass-through bacilli or those arising from colonization of the intestines. Serology provides a rapid and cost-effective alternative diagnostic tool and ELISA has high sensitivity in serological tests. Previous studies compared native semi-purified antigen (NSPA) of MAP prepared from a native strain of MAP of the 'Indian Bison type' genotype with commercial purified protoplasmic antigen (CPPA) of Allied Monitor Inc., (US) of 'cattle type' genotype in indirect (unabsorbed) ELISA kit and commercial (absorbed) ELISA kit for the screening of goats and sheep against MAP infection. This study reported better performance of the 'indigenous ELISA kit' (Singh *et al.*, 2007). Commercial ELISA kits showed lowered sensitivity

\*Corresponding author:

Email: [shoorvir\\_singh@rediffmail.com](mailto:shoorvir_singh@rediffmail.com).

and higher specificity in comparison to 'Indigenous ELISA kits' (Singh *et al.*, 2007). In the present study 'indigenous ELISA kit' initially developed for the screening of goats and sheep (Singh *et al.*, 2007) was modified (Singh *et al.*, 2009) and used for the screening of domestic livestock in north India.

## 2. Materials and Methods

### 2.1 Serum samples

A total of 829 serum samples were collected from different parts of North India between 2005 and 2008 and were screened at the Microbiology laboratory of Central Institute for Research on Goats (CIRG), Makhdoom, (UP). Of the 829 serum samples, 193, 143, 351 and 98 belonged to cattle, buffaloes, goats and sheep, respectively. Serum samples were collected aseptically, transported to the laboratory under ice and kept without adding any preservative at -20°C until used.

### 2.2 Control (negative and positive serum) samples

Serum samples collected from suspected, extremely weak cattle with known positive status for MAP in fecal culture were used as a positive control. Serum samples from healthy cattle belonging to the farmer's herd, with known culture-negative status for MAP infection, were used as a negative control for the screening of cattle and buffaloes serum samples. However, for goats and sheep positive and negative serum controls were selected from goats positive and negative in culture in earlier studies (Singh *et al.*, 2009).

### 2.3 Modified 'Indigenous absorbed ELISA kit'

'Indigenous ELISA kit' was used for the screening of MAP infection in domestic animals (Singh *et al.*, 2008) and was modified to improve the specificity by adding an absorption step using *M. phlei* culture (Allied Monitor Inc., USA), in the protocol (Singh *et al.*, 2009) and was used in the present study for the screening of domestic livestock.

### 2.4 Test proper

The 0.1µg of sonicated protoplasmic antigen in 100µl of carbonate-bicarbonate buffer (pH 9.6) was used for coating in duplicate wells of the flat bottom 96 well ELISA plate (Greiner Bio-One). The plates were kept overnight at 4°C followed by washing thrice with washing buffer PBST (PBS with 0.05% Tween 20). Blocking was done with 200µl of 3% skimmed milk in PBS and incubated at 37°C for one hour. After incubation plates were washed thrice with PBST, after washing the plates, 100µl of 1:50 diluted test serum samples in dilution buffer (PBST containing 1% BSA and 2mg/ml *M. phlei*) were added to each well, in the duplicates and incubated for 2 hours at 37°C. After

incubation three washings (5 minutes each) were given with PBST. Then 100µl of optimally diluted (1:8000) conjugate (Sigma) in PBS was added in all and incubated for an hour at 37°C. After incubation, the plate was washed three times with PBST. Finally, 200µl of a freshly prepared substrate (orthophenylene diamine dihydrochloride-OPD), 5mg per plate in substrate buffer (pH 5.0) was added to each well. Following incubation (in the dark) for 20 minutes at room temperature, absorbance was read at 450nm in an ELISA reader (Multiscan, Lab system, USA) without adding of stop solution (5N H<sub>2</sub>SO<sub>4</sub>). Blank, positive and conjugate controls were also run along with serum samples on each plate.

### 2.5 Calculation of cutoff

The cutoff value was calculated as per the method of Collins (2002). Serum samples were run in duplicate and mean OD was taken as the final OD value. OD values of the test samples were transformed to S/P ratio and animals in the strong positive category were considered positive for infection with MAP.

## 3. Results and Discussion

In the present study, using an 'indigenous absorbed ELISA' kit, the seroprevalence of MAP in domestic animals of North India was 22.1% (Table 1). A previous study from north India reported a slightly high (29.0%) prevalence of MAP in large ruminants using an indigenous unabsorbed ELISA kit as the diagnostic test (Singh *et al.*, 2008a). The lower prevalence of MAP in the present study may be due to the use of absorption of serum with *M. phlei*, which minimizes the cross-reaction of MAP with other mycobacteria (Yokomizo *et al.*, 1983). Though, cross-reactivity is the major problem for the diagnosis of mycobacterial infections (Yokomizo *et al.*, 1983). Species-wise 21.7, 16.0, 26.2 and 19.0% of samples were positive for MAP from cattle, buffalo, goat and sheep, respectively. The prevalence of MAP was higher in large ruminants (cattle and buffalo) in comparison to small ruminants (Goats and sheep). The higher prevalence of MAP in large ruminants may be attributed to the high endemicity of MAP infection and other production stress factors. The prevalence of MAP was higher in cattle (21.7%) as compared to buffaloes (16.0%) in north India. This finding indicated that the unproductivity of cattle in India may be to the JD. Similar to the present study, Singh *et al.*, (2008a), reported, 29.8 and 28.6% of animals were positive from cattle and buffaloes in North India using an unobserved ELISA test. Sivakumar *et al.*, (2005) screened the cattle sera from dairy and organized farms from Chennai, 21.3% were positive in ELISA and 10.8% of these ELISA-positive animals were confirmed by AGID. In another study, using ELISA kits in serum samples of

Buffaloes gave 46.7% seroprevalence (48.0% by target tissue culture) in farmer's buffaloes sacrificed for meat purposes in the Agra region (Yadav *et al.*, 2008). Siva Kumar *et al.*, (2005) reported 22% and a 10% incidence of MAP in cattle and buffaloes of Bareilly region.

As seen in large ruminants, a higher prevalence of MAP infection was found in small ruminants (goats and sheep). The prevalence of MAP was higher in goats (23.9%) as compared to sheep (19.0%). The higher prevalence of MAP in goats may be due to the high population density and endemicity of MAP in goat herds (Singh *et al.*, (1996); Singh *et al.*, (2008a); Vohra *et al.*, (2008). Similarly, Singh *et al.*, (1996) also reported a higher prevalence of MAP in goats as compared to sheep using microscopy and fecal culture. Recently, Singh *et al.*, (2008b) screened 35 goats from the Agra region by fecal culture and ELISA test, 77.1 and 40.0% of goats were positive for MAP infection, respectively. Of the 40% ELISA-positive goats, 71.4% were also positive in fecal culture. Previous studies also reported a higher prevalence of MAP in farm herds as

compared to farmer's herds (Kumar *et al.*, 2007). However, in the present study, no major difference was found in the prevalence of MAP in farm and farmer's herds. This may be due to the sharing of domestic livestock by the farm and farmer's herds in north India, the confinement of animals due to lack of pasture land and free grazing resources.

#### 4. Conclusion

The high prevalence of MAP infection in farm and farmer's herds of domestic ruminants in North India is directly related to the low per-animal productivity and there is an increased risk of exposure of human beings to MAP infection. Therefore, it is necessary to control MAP infection in animals first to safeguard the human population. National JD control programs are essential to reduce the prevalence of MAP in India and to improve per-animal productivity and also to reduce the risk of exposure to human beings.

**Table 1: Screening of domestic livestock against MAP infection by indigenous ELISA test.**

Ruminant	Species	Region	Management	Sample	Positives	Positives %
Large Ruminant	Cattle	Mathura	Farm	143	37	25.8
		Rohtak, Haryana	Farmer	24	7	29.1
		Bareilly	Farm	26	8	30.7
		Total			193	52
	Buffalo	Baghpat	Farm	38	7	18.4
		Rohtak, Haryana	Farmer	90	18	20.0
		Bareilly	Farm	15	4	26.6
Total			143	29	20.2	
Subtotal				336	81	24.1
Small Ruminant	Goat	CIRG, Mathura	Farm	163	32	19.6
		Etawah	Farm	80	22	37.5
		Agra	Farmer	35	14	40.0
		Ajmer	Farmer	73	16	21.9
	Total			351	84	23.9
	Sheep	CIRG, Mathura	Farm	43	11	25.5
		Mannavanur	Farm	98	16	16.3
Total			142	27	19.0	
Subtotal				493	111	22.5
<b>Total</b>				<b>829</b>	<b>192</b>	<b>23.1</b>

#### References

- [1]. Barbaruah, M.I. & Joseph, A.K. (2008). India: Dairy giant walking barefoot. Proceedings of the FAO-RAP Technical Meeting on Designing Effective Country-Specific Dairy Strategies, November 17-20, 2008, FAO, Rome, Italy.
- [2]. Chiodini, R.J., Van Kruiningen, H.J. & Merkal, R.S. (1984). Ruminant paratuberculosis (Johne's disease): the current status and future prospects. *Cornell Vet.*, 74(3): 218–262.
- [3]. Collins M.T. (2002). Interpretation of a commercial bovine paratuberculosis enzyme-linked immunosorbent assay by using likelihood ratios. *Clin. Diagn. Lab. Immunol.*, 9(6): 1367–1371. <https://doi.org/10.1128/cdli.9.6.1367-1371.2002>.
- [4]. Hasonova, L. & Pavlik, I. (2006). Economic impact of paratuberculosis in dairy cattle herds: a

- review. *Vet. Med.*, 51: 193-211. <https://doi.org/10.17221/5539-VETMED>.
- [5]. Kumar, P., Singh, S.V., Bhatiya, A.K., Sevilla, I., Singh, A.V., Whittington, R.J., Juste, R.A., Gupta, V.K., Singh, P.K., Sohal, J.S. & Vihan, V.S. (2007). Juvenile Capri-Paratuberculosis (JCP) in India: Incidence and characterization by six diagnostic tests. *Small Rumin. Res.*, 73: 45–53. <http://dx.doi.org/10.1016/j.smallrumres.2006.10.023>.
- [6]. Rideout, B.A., Brown, S.T., Davis, W.C., Gay, J.M., Giannella, R.A., Hines, M.E., Heuston, W.D. & Hutchinson, L.J. (2003). *Diagnosis and Control of Johne's Disease*. The National Academies Press, Washington, D.C.
- [7]. Sharma, G., Singh, S.V., Sevilla, I., Singh, A.V., Whittington, R.J., Juste, R.A., Kumar, S., Gupta, V.K., Singh, P.K., Sohal, J.S. & Vihan, V.S. (2008). Evaluation of indigenous milk ELISA with m-culture and m-PCR for the diagnosis of bovine Johne's disease (BJD) in lactating Indian dairy cattle. *Res. Vet. Sci.*, 84(1): 30–37. <https://doi.org/10.1016/j.rvsc.2007.03.014>.
- [8]. Singh, A.V., Singh, S.V., Sohal, J.S. & Singh, P.K. (2009). Comparative potential of modified indigenous, indigenous and commercial ELISA kits for diagnosis of *Mycobacterium avium* subspecies *paratuberculosis* in goat and sheep. *Indian J. Exp. Biol.*, 47(5): 379–382.
- [9]. Singh, N., Singh, S.V., Gupta, V.K., Sharma, V.D., Sharma, R.K. & Katoch, V.M. (1996). Isolation and identification of *Mycobacterium paratuberculosis* from naturally infected goatherds in India. *Indian J. Vet. Pathol.*, 20: 104-108.
- [10]. Singh, S.V., Singh, A.V., Singh, P.K. & Sohal, J.S. (2008b). An outbreak of caprine Paratuberculosis in a newly established commercial goat farm. *Haryana Vet.*, 47: 110-112.
- [11]. Singh, S.V., Singh, A.V., Singh, P.K., Sohal, J.S. & Singh, N.P. (2007). Evaluation of an indigenous ELISA for diagnosis of Johne's disease and its comparison with commercial kits. *Indian J. Microbiol.*, 47(3): 251–258. <https://doi.org/10.1007/s12088-007-0046-2>.
- [12]. Singh, S.V., Singh, A.V., Singh, R., Sharma, S., Shukla, N., Misra, S., Singh, P.K., Sohal, J.S., Kumar, H., Patil, P.K., Misra, P. & Sandhu, K.S. (2008a). Sero-prevalence of bovine Johne's disease in buffaloes and cattle population of North India using indigenous ELISA kit based on native *Mycobacterium avium* subspecies *paratuberculosis* 'Bison type' genotype of goat origin. *Comp. Immunol. Microbiol. Infect. Dis.*, 31(5): 419–433. <https://doi.org/10.1016/j.cimid.2007.06.002>.
- [13]. Sivakumar, P., Tripathi, B.N. & Singh, N. (2005). Detection of *Mycobacterium avium* subsp. *paratuberculosis* in intestinal and lymph node tissues of water buffaloes (*Bubalus bubalis*) by PCR and bacterial culture. *Vet. Microbiol.*, 108: 263–270. <https://doi.org/10.1016/j.vetmic.2005.04.002>.
- [14]. Vohra, J., Singh, S.V., Singh, A.V., Singh, P.K. & Sohal, J.S. (2008). Comparative distribution of *Mycobacterium avium* subspecies *paratuberculosis* in the target and non-target tissues of goats and sheep population in India. *Indian J. Anim. Sci.*, 78(4): 342–347.
- [15]. Whitlock, R.H. & Buergelt, C. (1996). Preclinical and clinical manifestations of paratuberculosis (including pathology). *Vet. Clin. North Am. Food Anim. Pract.*, 12(2): 345–356. [https://doi.org/10.1016/s0749-0720\(15\)30410-2](https://doi.org/10.1016/s0749-0720(15)30410-2).
- [16]. Yadav, D., Singh, S.V., Singh, A.V., Sevilla, I., Juste, R.A., Singh, P.K. & Sohal, J.S. (2008). Pathogenic 'Bison-type' *Mycobacterium avium* subspecies *paratuberculosis* genotype characterized from riverine buffalo (*Bubalus bubalis*) in North India. *Comp. Immunol. Microbiol. Infect. Dis.*, 31(4): 373–387. <https://doi.org/10.1016/j.cimid.2007.06.007>.
- [17]. Yokomizo, Y., Merkal, R.S. & Lyle, P.A. (1983). Enzyme-linked immunosorbent assay for detection of bovine immunoglobulin G1 antibody to a protoplasmic antigen of *Mycobacterium paratuberculosis*. *Am. J. Vet. Res.*, 44(11): 2205–2207.