



Leaf Area, Fresh Weight and Dry Weight Prediction Models for Ornamental Plants *Ficus benjamina* (cv. Starlight)

F. Bidarnamani*, H. Zarei, K. Mashayekhi and B. Kamkar

*Gorgan University of Agricultural Sciences and Natural Resources, Golestan Province, Gorgan, I.R., Iran.

Abstract: Measurements of leaf growth indices namely leaf area, fresh weight and dry weight are of value in physiological studies and plant growth estimation. The use of prediction models to estimate leaf area, fresh weight and dry weight is simple, rapid and nondestructive. Several mathematical functions have been formulated for estimating leaf area, fresh weight and dry weight of various crops but almost there is no information for *Ficus benjamina*. This work was aimed to propose leaf area (LA), fresh weight (FW) and dry weight (DW) prediction models for *Ficus benjamina* (cv. Starlight) leafy ornamental pot plant using leaf length (L) and width (W). 1000 leaves were collected randomly from greenhouse ornamental plants and 700 of cuts were used for prediction models. LA was measured with a digital area meter (DELTA-T, Co. Durham, UK), related FW and DW also were weighted and leaf dimensions were determined by the ruler. For each studying growth index LA, FW and DW the predictive abilities of three regression equations (linear, polynomial and power) were compared with different independent variables for each equation. Leaf length \times width provided a good estimation of leaf area and fresh weight of the leaves of *Ficus benjamina*. It was also concluded that leaves the dry weight of *Ficus benjamina* can be estimated or simulated as a power function of $L \times W$ or $L + W$ with reasonable accuracy. Moreover, a reasonable relationship between leaf fresh weight and leaf area was found too.

Keywords: Foliage pot plants, Leaf area estimation, Leaf growth estimation, Non-destructive methods, *Ficus benjamina*.

1. Introduction

Leaf growth indices including leaf area (LA), leaf fresh weight (FW), and leaf dry weight (DW) are important factors required in most physiological and agronomic studies involving plant growth (Guo and Sun, 2001). LA has a remarkable effect on growth and its estimation is a key component of crop growth models (Lizaso *et al.*, 2003; Roupael *et al.*, 2006). Different methods to measure leaf area (Evans, 1972; Causton & Venus, 1981; Norman & Campbell, 1991; Bignami and Rossini, 1996) are time-consuming, destructive and the facilities are commonly expensive. Therefore, it is of value to find a non-destructive and repeatable method during the growth period which is inexpensive and reduces the experimental variability associated to destructive sampling procedures (NeSmith, 1992; De Swart *et al.*, 2004). Several

prediction models have been determined to estimate leaf area for numerous horticultural crops (Potdar and Pawar, 1991; Montero *et al.*, 2000; Stoppani *et al.*, 2003; Demirsoy *et al.*, 2004; De Swart *et al.*, 2004; Salerno *et al.*, 2005; Roupael *et al.*, 2006; Serdar and Demirsoy, 2006; Cho *et al.*, 2007; Cristofori *et al.*, 2007; Mendoza-de Gyves *et al.*, 2007; Peksen, 2007; Rivera *et al.*, 2007; Tsialtas *et al.*, 2008; Olfati *et al.*, 2009; Roupael *et al.*, 2010). In a number of research works, there is a strong relationship between L and W and LA, FW and DW (Mokhtarpour *et al.*, 2010) or DW (Karimi *et al.*, 2010) appeared to be reasonably related to leaf area. While information on the estimation of ornamental plant, LA, FW and DW, especially *Ficus benjamina* is still deficient. The objective of this study was focused to develop non-destructive models for estimating LA, FW and DW of *Ficus benjamina*, based on leaf length and width.

*Corresponding author:

E-mail: f.bidarnamani@yahoo.com, hosszareei@yahoo.co.uk.

2. Materials and Methods

2.1 Model construction

The experiment was conducted in north of Iran (Gorgan; Latitude: N 37°00' to 37°30' and Longitude: E 54°00' to 54°30'). The eco-climate of Gorgan is moderate and humid. The present investigation was carried out on small pot plants raised from semi hardwood cuttings transplanted in 4 liter plastic pots containing 22 different pot mixtures in glass-house conditions. Sampling was done at 9 months after beginning of growth in spring and a total of 1000 leaves harvested for this experiment. The length from the lamina tip to the intersection of the lamina and petiole along the lamina midrib and the maximum width of all leaves were measured manually with a simple ruler to the nearest 1.0mm (Fig. 1). The leaf area measured with a digital leaf area meter (DELTA-T, Co. Durham, UK) to the nearest 1.0cm². Fresh weight of each leaf also was measured and then dry weights were determined after drying in 70°C oven for 48 hours. The fresh and dry weight of leaves was measured to the nearest 0.001g. The relationships between LA, FW and DW as a dependent variable and length (L), width (W), L+W, L×W, L² and W² as independent variables and also LA as dependent variable and FW and DW as independent variable were determined using regression analysis on a total data of 700 leaves. Coefficients of determination (R²) were calculated and equations with the highest R² were used in the final estimations. The linear, polynomial and power functions were developed through SAS software (SAS Institute, 1992) and Excel worksheet.




Fig. 1. *Ficus benjamina* leaves showing the point of leaf length (L) and width (W) measurement.

The estimated LA, FW, and DW were determined by fitting the equations. Then estimated and measured LA, FW and DW were compared with testing the significance of regression equation and degree of goodness of fit R² between estimated and observed values. The model with lower root mean square error (RMSE), lower relative mean absolute error (RMAE), higher R², lower bias of the linear regressed line between observed versus predicted values from the 1:1 line and lower coefficient of variance (CV) was

selected as the best model to estimate leaf area by following formulas.

$$\text{Bias} = \frac{1}{N} \sum_{i=1}^N D_i \quad (1)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (D_i)^2} \quad (2)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (D_i)^2} \quad (3)$$

$$\text{CV} = \frac{\text{RMSE}}{\bar{Y}} \times 100 \quad (4)$$

Where, N is the total number of situations, Y_i and \hat{Y} are the observed and predicted Y values, respectively, and $D_i = Y_i - \hat{Y}$ (Wallach, 2006).

'a' and 'b' (as intercept and slope values of linear regression between observed versus predicted values of leaf area) were compared with zero and 1. A closer 'a' to zero and closer 'b' to 1 indicate better estimates of models.

2.2 Model validation

In order to validate the selected models for estimating LA, FW and DW about 300 leaves were randomly sampled and LA, L and W were determined by the previously described procedures. These values were used as independent data to validate the models (were not used in model fitting). The slope and intercept of the model were tested to see if they were significantly different from the slope and intercept of the 1:1 correspondence line. Regression analyses were conducted using SAS software (SAS Institute, 1992).

3. Result

Various mathematical models for indirect estimation of leaf area and lesser leaf fresh weight and leaf dry weight of different plant species have been described. The outcomes of the present study were in agreement with some of the previous studies mentioned above on non-destructive model development for predicting leaf area, leaf fresh weight and leaf dry weight using simple linear leaf measurements. Different prediction equations obtained for estimating the LA, FW and DW of *Ficus benjamina* (cv. Starlight) involving different independent variables, viz, L, W, L², W², L+W and L×W were formulated for estimating leaf area by using different equations. The equations with lower R² and higher CV were eliminated at the beginning of this study (Fig. 2).

However, for estimating LA by L and W, linear and polynomial functions using a product of L and W viz., Eq. (6) and (5) [$Y = a + b(L \times W) + c(L \times W)^2$ and $Y = a + b(L \times W)$ respectively] had a higher R² value (0.950 and 0.949) with a lower bias (-0.0004 and -0.0005), lower RMSE (0.74 and 0.75), lower RMAE (0.090 and 0.092) and lower CV (10.20% and 10.30%) than other

equations tested and showed the best LA estimation (Table 1, Fig. 2: no. 1 and 2). Evaluation of these two models by 300 data from independent leaf length and width measurements showed that both models are efficient to estimate LA (Table 2, Fig. 3: no. 1 and 2).

For fresh weight, power function viz., Eq. (4) Using the product of length and width described the good relationship which had approximately higher R² value (0.838), lower bias (0.002), lowest RMSE (0.029), lowest RMAE (0.137) and lower CV (16.94%) than other equations tested and showed the best FW estimation (Table 3, Fig. 2: no. 3). Evaluation of this model by 300 data from independent leaf length and

width measurements showed that this model is practical to estimate FW (Table 4, Fig. 3: no. 3).

Using the same method as in LA and FW for estimating dry weight by measuring length and width, power functions viz., Eq. (4) and (3) using correspondingly product and sum of leaf length and width showed the highest R² value (0.771 and 0.761), nearly lower bias (0.0007 and -0.0014), lowest RMSE among the other models to estimate DW (Table 5, Fig. 2: no. 4 and 5). Validation of these two models by 300 data from independent leaf length and width measurements showed that both models are appropriate to estimate FW (Table 6, Fig. 3: no. 4 and 5).

Table 1. Slope (b) and intercept (a) values of the models used to estimate the *Ficus benjamina* (cv. Starlight) leaf area (LA) of single leaves from length (L) and width (W) measurements.

Model number	Regression model	R ²	Bias	RMSE	RMAE	CV	a	b
1	LA = 0.3248L ² - 0.5838L - 1.0012	0.886	-0.00100	1.120	0.140	15.38	0.829	0.896
2	LA = 0.8983W ^{2.1271}	0.903	0.06900	1.050	0.119	14.45	0.591	0.909
3	LA = 1.078W ² - 0.0255W - 0.2526	0.900	0.00060	1.050	0.120	14.40	0.726	0.900
4	LA = 0.1562(L+W) ² - 0.4731(L+W) - 0.9684	0.941	0.00030	0.810	0.101	11.07	0.429	0.941
5	LA = 0.573(L×W) - 0.8412	0.949	-0.00050	0.750	0.092	10.30	0.371	0.949
6	LA = 0.0025(L×W) ² + 0.4956(L×W) - 0.3263	0.950	-0.00040	0.740	0.090	10.20	0.364	0.950
7	LA = 0.2703(L ²) - 0.4971	0.886	-0.00110	1.124	0.138	15.41	0.886	0.833
8	LA = 0.0006(L ²) + 0.2331(L ²) - 0.0065	0.887	-0.01100	1.120	0.140	15.37	0.821	0.889
9	LA = 0.8983(W ²) ^{1.0635}	0.903	0.07000	1.054	0.119	14.46	0.591	0.909
10	LA = -0.0011(W ²) ² + 1.0902(W ²) - 0.341	0.900	0.00040	1.049	0.120	14.40	0.726	0.900
11	LA = 1.0732(W ²) - 0.2848	0.900	-0.00007	1.050	0.120	14.40	0.726	0.900

Table 2. Validation of models for estimation of total leaf area of *Ficus benjamina* cuttings.

Model number	R ²	Bias	RMSE	RMAE	CV	a	b
1	0.893	-0.124	0.891	0.130	13.73	0.798	0.896
2	0.810	-0.037	1.215	0.154	18.72	0.602	0.913
3	0.810	-0.118	1.217	0.154	18.74	0.721	0.907
4	0.921	-0.079	0.767	0.106	11.81	0.409	0.949
5	0.913	-0.089	0.814	0.106	12.55	0.325	0.964
6	0.914	-0.077	0.804	0.106	12.38	0.349	0.958
7	0.891	-0.138	0.901	0.129	13.88	0.781	0.901
8	0.893	-0.132	0.892	0.130	13.75	0.800	0.897
9	0.810	-0.037	1.215	0.154	18.72	0.602	0.913
10	0.810	-0.119	1.218	0.154	18.76	0.717	0.908
11	0.810	-0.119	1.217	0.154	18.74	0.721	0.907

Table 3. Slope (b) and intercept (a) values of the models used to estimate the *Ficus benjamina* (cv. Starlight) leaf fresh weight (FW) of single leaves from length (L) and width (W) measurements.

Model number	Regression model	R ²	Bias	RMSE	RMAE	CV	a	b
1	FW = 0.0067L ^{1.8617}	0.762	0.004	0.034	0.164	20.10	0.044	0.714
2	FW = 0.0246W ^{1.8779}	0.810	0.003	0.031	0.144	18.15	0.035	0.778
3	FW = 0.0025(L+W) ^{1.9813}	0.824	0.001	0.030	0.144	17.51	0.035	0.791
4	FW = 0.0109(L×W) ^{0.997}	0.838	0.002	0.029	0.137	16.94	0.032	0.803
5	FW = 0.0067(L ²) ^{0.9308}	0.762	0.004	0.034	0.164	20.10	0.044	0.714
6	FW = 0.0246(W ²) ^{0.9389}	0.810	0.003	0.031	0.144	18.15	0.035	0.778

FW, leaf fresh weight; L, leaf length; W, leaf width

Table 4. Validation of models for estimation of total leaf area of *Ficus benjamina* cuttings.

Model number	R ²	Bias	RMSE	RMAE	CV	a	b
1	0.718	0.006	0.028	0.148	18.76	0.036	0.719
2	0.736	0.006	0.028	0.134	18.50	0.020	0.828
3	0.779	0.005	0.025	0.126	16.57	0.024	0.806
4	0.790	0.006	0.024	0.123	16.37	0.020	0.825
5	0.718	0.006	0.028	0.148	18.76	0.036	0.719
6	0.736	0.006	0.028	0.134	18.50	0.020	0.828




Fig. 2. Selected regression models for estimating the leaf area, fresh and dry weight of *Ficus benjamina* from the leaf length by width product ($L \times W$) and sum ($L + W$) and also estimating the leaf area from fresh weight.

Table 5. Slope (b) and intercept (a) values of the models used to estimate the *Ficus benjamina* (cv. Starlight) leaf dry weight (DW) of single leaves from length (L) and width (W) measurements.

Model number	Regression model	R ²	Bias	RMSE	RMAE	CV	a	b
1	$DW = 0.0019L^{1.9176}$	0.7057	0.0008	0.0120	0.188	22.69	0.063	0.6761
2	$DW = 0.0065W^2 - 0.0002W + 0.0031$	0.7172	0.0001	0.0113	0.178	21.26	0.0179	0.666
3	$DW = 0.0007(L+W)^{2.0437}$	0.7609	-0.0014	0.011	0.177	20.83	0.0159	0.6578
4	$DW = 0.0031(L \times W)^{1.0278}$	0.7709	0.0007	0.0107	0.158	20.22	0.0168	0.6786
5	$DW = 0.0001(W^2)^2 + 0.004(W^2) + 0.0144$	0.736	0.0014	0.0114	0.170	21.43	0.6266	0.0189
6	$DW = 0.0019(L^2)^{0.9588}$	0.7057	0.0008	0.0120	0.188	22.69	0.0187	0.6421

DW, leaf dry weight; L, leaf length; W, leaf width

Table 6. Validation of models for estimation of total leaf area of *Ficus benjamina* cuttings.

Model number	R ²	Bias	RMSE	RMAE	CV	a	b
1	0.737	0.0019	0.008	0.130	17.52	0.008	0.782
2	0.745	0.0012	0.008	0.126	17.59	0.004	0.876
3	0.795	0.0006	0.007	0.118	15.44	0.004	0.895
4	0.810	0.0024	0.007	0.119	15.86	0.006	0.906
5	0.746	0.0025	0.008	0.133	18.19	0.003	0.877
6	0.737	0.0019	0.008	0.130	17.52	0.008	0.782




Fig. 3. Comparison of predicted and observed areas, fresh weight and dry weight of *Ficus benjamina* leaves using selected models. [FW-fresh weight; L-length; W-width].

Table 7. Slope (b) and intercept (a) values of the models used to estimate the *Ficus benjamina* (cv. Starlight) Leaf Area (LA) of single leaves from leaf fresh weight (F) and leaf dry weight (D) measurements.

Model number	Regression model	R ²	Bias	RMSE	RMAE	CV	a	b
1	$LA = 45.611F^{0.9798}$	0.850	0.0870	1.333	0.138	16.50	1.055	0.859
2	$LA = -280.42D^2 + 160.68D + 0.4444$	0.719	0.0007	1.700	2.890	21.04	2.274	0.719
3	$LA = 127.98D + 1.2646$	0.719	0.0003	1.707	0.195	21.12	2.291	0.716
4	$LA = -15.111(F+D)^2 + 41.797(F+D) - 0.3654$	0.834	0.0004	1.307	0.146	16.18	1.345	0.834
5	$LA = 34.465(F+D) + 0.4093$	0.832	-0.0004	1.314	0.147	16.26	1.358	0.832
6	$LA = 78.228(F \times D)^{0.4838}$	0.813	0.1120	1.362	0.150	16.87	1.213	0.836
7	$LA = -8041.8(F \times D)^2 + 608.8(F \times D) + 3.149$	0.811	0.0066	1.394	0.166	17.25	1.528	0.811
8	$LA = 45.611(F^2)^{0.4899}$	0.850	0.0870	1.331	0.138	16.48	1.055	0.859
9	$LA = -67366(D^2)^2 + 1693.6(D^2) + 3.674$	0.704	-0.0004	1.734	0.208	21.53	2.389	0.704

LA, Leaf area; F, leaf fresh weight; D, leaf dry weight

Table 8. Validation of models for estimation of total leaf area *Ficus benjamina* cuttings.

Model number	R ²	Bias	RMSE	RMAE	CV	a	b
1	0.8209	-0.681	1.316	0.181	19.87	0.7474	0.99
2	0.8191	-0.635	1.219	0.177	18.40	1.5794	0.8573
3	0.8181	-0.624	1.212	0.172	18.30	1.7381	0.8319
4	0.8478	-0.727	1.245	0.176	18.80	0.8673	0.9788
5	0.8487	-0.720	1.225	0.175	18.50	0.9859	0.9598
6	0.8601	-0.540	1.093	0.151	16.50	0.7803	0.9637
7	0.8545	-0.665	1.159	0.174	17.50	1.1295	0.9298
8	0.8209	-0.681	1.316	0.181	19.87	0.7474	0.99
9	0.8066	-0.621	1.238	0.192	18.69	1.8534	0.8139

To estimate leaf area by measuring leaf fresh and dry weight, we obtained power functions viz., Eq. (8) & (1) using a fresh weight which had well relationships with LA, cleared in high R² (0.850), nearly lowest RMSE (1.331 & 1.333), lowest RMAE (0.138) and nearly lowest CV (16.48 & 16.50) values among the other equations (Table 7, Fig. 3: no. 6 & 7). Comparisons were made between measured versus calculated leaf area of other 300 leaves that were not involved in modeling set by using selected models. It was found that the Eq. (1) may be better than Eq. (8) to estimate leaf area using fresh weight measurement (Table 8, Fig. 3: no. 6 and 7).

4. Discussion

Several studies have been carried out to estimate leaf area by measuring leaf length and width (Robbins and Pharr, 1987; Gamiely *et al.*, 1991; Montero *et al.*, 2000; Williams III and Martinson, 2003). We found that regression equation were fitted using the length and the width of the leaves, the product of them and their squares as independent variables to estimate leaf area, fresh and dry weights. Leaf length or width exclusively did not offer a good variable to estimate leaf growth indices, and it is in similarity with Karimi *et al.*, (2009) but indifference with Williams Iii and Martinson, (2003) and Cho *et al.*, (2007) that declared a solitary variable of either leaf length or leaf width has a good correspondence with leaf area and or leaf weight.

Our research results indicated that the product of leaf length and width is a proper variable to verify LA and FW and both product and sum of L and W are suitable variables to determine DW using regression models. The variable with the highest explanatory capability was used to develop a general equation to predict LA, FW and DW. It was revealed that areas of this type of *Ficus* leaves are well correlated to the product of its length and width with high R² values. R² values for fresh weight (0.838) and dry weight (0.771) of leaves were less than the R² value of leaf area (0.950), as Cho *et al.*, (2007) and Krimi *et al.*, (2009) found and this was reasoned by Cho *et al.*, (2007) that it is probably due to differences in specific leaf area. They stated that SPAD data are useful to determine leaf yield; but in this study, SPAD was not investigated.

There are lots of works which suggest calculation of leaf area from leaf dry weight data (Sharrett and Baker, 1985; Ma *et al.*, 1992; Akram-Ghaderi and Soltani, 2007) but we found a reasonable relationship between leaf fresh weight (not dry weight) and leaf area. Power function described this relationship better than the other equation types.

Validation of selected models by independent data showed a strong conformity between predicted and measured data. According to these results, leaf length and width contribute to rapid, simple and non-destructive determination leaf area, leaf fresh and dry weight. Moreover, leaf fresh and dry weight could be used to estimate leaf area of *Ficus benjamina* (cv. Starlight). Further studies are required to evaluate these models or constructing new models for other cultivars of this ornamental plant.

References

- [1]. Akram-ghaderi, F. & Soltani, A. (2007). Leaf area relationships to plant vegetative characteristics in cotton (*Gossypium hirsutum* L.) grown in a temperate subhumid environment. International Journal of Plant Production, 1: 63-71. SAS Institute, 1992. SAS/STAT user's guide, Version 6, 4th editions, SAS Inst., Inc., Cary, NC.
- [2]. Bignami, C. & Rossini, F. (1996). Image analysis, estimation of leaf area index and plant size of young hazelnut plants. *J. Hort. Sci.*, 71: 113-121.
- [3]. Causton, D.R. & Venus, J.C. (1981). The biometry of plant growth. London: Edward Arnold, 301p.
- [4]. Cho, Y.Y., Oh, S., Oh, M.M. & Son, J.E. (2007). Estimation of individual leaf area, fresh weight, and dry weight of hydroponically grown cucumbers (*Cucumis sativus* L.) using leaf length, width and SPAD value. *Sci. Hort.*, 111: 330-334.
- [5]. Cristofori, V., Roupheal, Y., Mendoza-de Gyves, E. & Bignami, C. (2007). A simple model for estimating leaf area of hazelnut from linear measurements. *Sci. Hort.*, 113: 221-225.
- [6]. De Swart, E.A.M., Groenwold, R., Kanne, H.J., Stam, P., Marcelis, L.F.M. & Voorrips, R.E. (2004). Non-destructive estimation of leaf area for different plant ages and accessions of *Capsicum annum* L. *J. Hort. Sci. Biotechnol.*, 79: 764-770.

- [7]. Demirsoy, H., Demirsoy, L., Uzun, S. & Ersoy, B. (2004). Non-destructive leaf area estimation in peach. *Eur. J. Hort. Sci.*, 69: 144-146.
- [8]. Evans, G.C. (1972). The quantitative analysis of plant growth. Oxford: Blackwell. 734p.
- [9]. Gamiely, S., Randle, W.M., Mills, H.A. & Smittle, D.A. (1991). A rapid and nondestructive method for estimating leaf area of onions. *Hort. Sci.*, 26(2): 206.
- [10]. Guo, D.P. & Sun, Y.Z. (2001). Estimation of leaf area of stem lettuce (*Lactuca sativa* var *angustana*) from linear measurements. *Indian Journal of Agricultural Science*, 71(7): 483-486.
- [11]. Karimi, S., Tavallali, V., Rahemi, M., Rostami, A.A. & Vaezpour, M. (2009). Estimation of leaf growth on the basis of measurements of leaf lengths and widths, choosing Pistachio seedling as model. *Aust. J. Basic & Appl. Sci.*, 3(2): 1070-1075.
- [12]. Lizaso, J.I., Batchelor, W.D. & Westgate, M.E. (2003). A leaf area model to simulate cultivar-specific expansion and senescence of maize leaves. *Field Crops Res.*, 80: 1-17.
- [13]. Ma, L., Gardner, F.P. & Selamat, A. (1992). Estimation of leaf area from leaf and total mass measurements in peanut. *Crop Science*, 32: 467-471.
- [14]. Mendoza-de Gyves, E., Rouphael, Y., Cristofori, V. & Rosana Mira, F. (2007). A non-destructive, simple and accurate model for estimating the individual leaf area of kiwi (*Actinidia deliciosa*). *Fruits*, 62: 171-176.
- [15]. Mokhtarpour, H., B.S. Teh C, Saleh, G., Selamat, A.B., Asadi, M.E. & Kamkar, B. (2010). Non-destructive estimation of the maize leaf area, fresh weight, and dry weight using leaf length and leaf width. *Communication in Biometry and Crop Science*, 5(1): 19-26.
- [16]. Montero, F.J., de Juan, J.A., Cuesta, A. & Brasa, A. (2000). Nondestructive methods to estimate leaf area in *Vitis vinifera* L. *Hort. Sci.*, 35(4): 696-698.
- [17]. NeSmith, D.S. (1992). Estimating summer squash leaf area nondestructively. *Hort. Sci.*, 27: 77.
- [18]. Norman, J.M. & Campbell, G.S. (1991). Canopy structure. In: Percy, R.W., Ehleringer, J.R., MOONEY, H.A., Rundel, P.W. (Ed.). *Plant physiological ecology: field methods and instrumentation*. London: Chapman & Hall., p.301-325.
- [19]. Olfati, J.A., Peyvast, Gh., Sanavi, M., Salehi, M., Mahdipour, M. & Nosratie-Rad, Z. (2009). Comparisons of leaf area estimation from linear measurements of red cabbage. *Int. J. Veg. Sci.*, 15: 185-192.
- [20]. Peksen, E. (2007). Non-destructive leaf area estimation model for faba bean (*Vicia faba* L.). *Sci. Hort.*, 113: 322-328.
- [21]. Potdar, M.V. & Pawar, K.R. (1991). Non-destructive leaf area estimation in banana. *Sci. Hort.*, 45, 251-254.
- [22]. Rivera, C.M., Rouphael, Y., Cardarelli, M. & Colla, G. (2007). A simple and accurate equation for estimating individual leaf area of eggplant from linear measurements. *Europ. J. Hort. Sci.*, 72: 228-230.
- [23]. Robbins, N.S. & Pharr, D.M. (1987). Leaf area prediction methods for cucumber from linear measurements. *Hort. Sci.*, 22 (6): 1264-1266.
- [24]. Rouphael, Y., Mouneimne, A.H., Rivera, C.M., Cardarelli, M., Marucci, A. & Colla, G. (2010). Allometric models for non-destructive leaf area estimation in grafted and un-grafted watermelon (*Citrullus lanatus* Thunb.). *J. Food Sci. Environ.*, 8: 161-165.
- [25]. Rouphael, Y., Rivera, C.M., Cardarelli, M., Fanasca, S. & Colla, G. (2006). Leaf area estimation from linear measurements in zucchini plants of different ages. *J. Hort. Sci. Biotechnol.*, 81: 238-241.
- [26]. Salerno, A., Rivera, C.M., Rouphael, Y., Colla, G., Cardarelli, M., Pierandrei, F., Rea, E. & Saccardo, F. (2005). Leaf area estimation of radish from linear measurements. *Adv. Hort. Sci.*, 19: 213-215.
- [27]. Serdar, U. & Demirsoy, H. (2006). Non-destructive leaf area estimation in Chestnut. *Sci. Hort.*, 108: 227-230.
- [28]. Sharrett, B.S. & D.G. Baker. (1985). Alfalfa leaf area as a function of dry matter. *Crop Science*, 26: 1040-1042.
- [29]. Stoppani, M.I., Wolf, R., Francescangeli, N. & Martí, H.R. (2003). A non-destructive and rapid method for estimating leaf area of broccoli. *Adv. Hort. Sci.*, 17: 173-175.
- [30]. Tsialtas, J.T., Koundouras, S. & Zioziou, E. (2008). Leaf area estimation by simple measurements and evaluation of leaf area prediction models in Cabernet-Sauvignon grapevine leaves. *Photosynthetica*, 46: 452-456.
- [31]. Williams III, L. & Martinson, T.E. (2003). Nondestructive leaf area estimation of 'Niagara' and 'De Chaunac' grapevines. *Sci. Hort.*, 98: 493-498.
- [32]. Wallach, D. (2006). Evaluating crop models. In D. Wallach, D. Makowski, & J.W. Jones, *Working with dynamic crop models* (pp. 11-19). Elsevier.